

Lagrangian statistics in fully developed turbulence

L. Biferale¹, G. Boffetta², A. Celani³, A. Lanotte⁴, and F. Toschi⁵

¹ *Dipartimento di Fisica and INFN, Università degli Studi di Roma “Tor Vergata”,
Via della Ricerca Scientifica 1, 00133 Roma, Italy*

² *Dipartimento di Fisica Generale and INFN, Università degli Studi di Torino, Via Pietro Giuria 1, 10125, Torino, Italy*

³ *CNRS, INLN, 1361 Route des Lucioles, 06560 Valbonne, France*

⁴ *CNR-ISAC, Str. Prov. Lecce-Monteroni km.1200, 73100 Lecce, Italy*

⁵ *Istituto per le Applicazioni del Calcolo, CNR, Viale del Policlinico 137, 00161 Roma, Italy*

(Dated: February 8, 2008)

The statistical properties of fluid particles transported by a fully developed turbulent flow are investigated by means of high resolution direct numerical simulations. Single trajectory statistics are investigated in a time range spanning more than three decades, from less than a tenth of the Kolmogorov timescale up to one large-eddy turnover time. Acceleration and velocity statistics show a neat quantitative agreement with recent experimental results. Trapping effects in vortex filaments give rise to enhanced small-scale intermittency on Lagrangian observables.

PACS numbers: 47.27.-i, 47.10.+g

The knowledge of the statistical properties of particle tracers advected by a fully developed turbulent flow is a key ingredient for the development of stochastic Lagrangian models in such diverse contexts as turbulent combustion, pollutant dispersion, cloud formation and industrial mixing [1, 2, 3, 4]. Given the importance of this problem, there are comparatively few experimental studies of turbulent Lagrangian dispersion. Progress in this direction has been hindered mainly by the presence of a wide range of dynamical timescales, an inherent property of fully developed turbulence. Indeed, in order to obtain an accurate description of the particle statistics it is necessary to follow their paths with very high resolution, i.e., well below the Kolmogorov timescale τ_η , and for a long time lapse – of the order of an eddy turnover time T_L . The ratio of these timescales can be estimated as $T_L/\tau_\eta \sim R_\lambda$ where the microscale Reynolds number R_λ ranges in the hundreds for typical laboratory experiments. A recent breakthrough has been made by La Porta *et al* [5, 6] who, borrowing techniques from high-energy physics, were able to track three-dimensional trajectories with a resolution of $0.05\tau_\eta$ and thus to study the statistics of particle acceleration. However, owing to the small measurement volume, trajectories could be followed only up to a few τ_η , preventing any investigation of the long time correlations along particle paths. Conversely, the acoustical technique adopted by Mor-dant *et al* [7] enabled them to successfully track particles for durations comparable to T_L but could not access time delays of the order of τ_η , and was restricted to one-dimensional measurements. More conventional techniques, e.g. based on CCD cameras [8], reveal useful only for moderate Reynolds numbers ($R_\lambda \simeq 100$) due to their limitations in the acquisition rate. In addition, all laboratory techniques require a considerable amount of signal processing to get rid of experimental noise. Besides these drawbacks, increasing difficulties are met when attempting to deal with multi-particle tracking.

In the light of the above remarks Direct Numerical Simulation of fully developed turbulence represents a valuable alternative tool for the investigation of Lagrangian statistics [9, 10, 11, 12, 13, 14, 15, 16, 17] and its important contribution has been recently reviewed in Ref. [4]. Nonetheless, computational approaches have to face three demanding requirements: (i) have the largest possible value of R_λ in order to maintain the flow in a fully developed turbulent state; (ii) resolve properly the dissipative scales with the aim of computing accurately small-scale observables; (iii) follow particle trajectories for times comparable to a large-eddy-turnover time. High resolution and huge computational resources are therefore needed to accomplish such a goal, making massive parallel computing by far and away the most appropriate tool. In this Letter we report the results of Direct Numerical Simulations of Lagrangian transport in homogeneous and isotropic turbulence on the parallel computer IBM-SP4 at CINECA, on 512^3 and 1024^3 cubic lattices with Reynolds numbers up to $R_\lambda \sim 280$, a very accurate resolution of dissipative scales, and a long integration time $T \approx T_L$. The Navier-Stokes equations

$$\partial_t \mathbf{u} + \mathbf{u} \cdot \nabla \mathbf{u} = -\nabla p + \nu \Delta \mathbf{u} , \quad (1)$$

are integrated on a triply periodic box by means of a fully dealiased pseudospectral code (the numerical parameters are listed in Table I). Energy is injected at an average rate ϵ by keeping constant the total energy in each of the first two wavenumber shells [18]. With the present choice of parameters the dissipative range of lengthscales is extremely well resolved. With regard to the scaling properties of the velocity field, our results are in perfect numerical agreement with previous numerical simulations at comparable R_λ (see e.g. [19]). Upon having reached statistically stationary conditions for the velocity field, millions of Lagrangian tracers have been seeded into the flow and their trajectories integrated according

R_λ	u_{rms}	ε	ν	η	L	T_L	τ_η	T	δx	N^3	N_p
183	1.5	0.886	0.00205	0.01	3.14	2.1	0.048	5	0.012	512^3	$0.96 \cdot 10^6$
284	1.7	0.81	0.00088	0.005	3.14	1.8	0.033	4.4	0.006	1024^3	$1.92 \cdot 10^6$

TABLE I: Parameters of the numerical simulations. Microscale Reynolds number R_λ , root-mean-square velocity u_{rms} , energy dissipation ε , viscosity ν , Kolmogorov lengthscale $\eta = (\nu^3/\varepsilon)^{1/4}$, integral scale L , large-eddy turnover time $T_L = L/u_{rms}$, Kolmogorov timescale $\tau_\eta = (\nu/\varepsilon)^{1/2}$, total integration time T , grid spacing δx , resolution N^3 , and the number of Lagrangian tracers N_p .




FIG. 1: Snapshots of a subset of Lagrangian particles (≈ 12000) at three different times $t = (0.2, 0.5, 0.8)T_L$. Gray scales encode different velocity amplitudes.

to

$$\frac{d\mathbf{X}}{dt} = \mathbf{u}(\mathbf{X}(t), t), \quad (2)$$

over a time lapse of the order of T_L (See Fig. 1). Particle positions $\mathbf{X}(t)$ and velocities $\mathbf{v}(t) = \dot{\mathbf{X}}(t)$ have been stored at a sampling rate $0.07\tau_\eta$. The forces acting on the particle – pressure gradients $\nabla p(\mathbf{X}(t), t)$, viscous forces $\nu\Delta\mathbf{u}(\mathbf{X}(t), t)$ and external input – and the resulting particle acceleration $\mathbf{a}(t) = \dot{\mathbf{v}}(t)$ have been recorded along the particle paths every $0.14\tau_\eta$. The resulting database permits detailed study of the statistics of Lagrangian velocity and acceleration over a range of timescales spanning more than three decades.

In what follows, we will focus on the description of the statistics of single particle trajectories, deferring the discussion of multi-particle statistics to a forthcoming publication. The analysis reveals the presence of frequent entrapment events within vortical structures (see Fig. 2). The characteristic frequency of these spiraling paths is comparable to τ_η^{-1} and their duration can be as long as $10\tau_\eta$. The velocity experienced during these events can attain $5u_{rms}$ with an ensuing acceleration $5u_{rms}/\tau_\eta$ as large as $80a_{rms}$. As we will show below, these events have a significant impact on time correlations up to $10\tau_\eta$.

In Fig. 3 we present the probability density functions (PDFs) of velocity differences $\delta_\tau v = v(t+\tau) - v(t)$ taken along a given direction and rescaled by their variance $\langle(\delta_\tau v)^2\rangle^{1/2}$. For the shortest time delays shown in the figure the velocity increment PDF approaches the distribution of the acceleration $a(t)$ which is characterized by a strong intermittency, stretched exponential tails and a




FIG. 2: Trajectory and time series. Left panel: 3D trajectory of a trapping event in vortex filament. Acceleration and velocity fluctuations here reach about 30 and 2 r.m.s. values, respectively (right panels).

flatness $\langle a^4 \rangle / \langle a^2 \rangle^2 \approx 40$. Classical Kolmogorov scaling arguments (see e.g. Ref. [20]) yield for the acceleration variance the prediction

$$\langle a^2 \rangle = a_0 \varepsilon^{3/2} \nu^{-1/2}. \quad (3)$$

We measure the constant $a_0 = 3.5 \pm 0.3$, in agreement with the results of Ref. [13] at comparable R_λ . Yet, one needs to notice that intermittent corrections may introduce in the definition of a_0 some dependency on Reynolds number[6]. At larger time separations the PDFs are decreasingly intermittent and eventually be-




FIG. 3: Probability density functions of velocity increments and acceleration, normalized with their variance. Curves refer to time increments $\tau = (97, 25, 6, 0.7)\tau_\eta$ from inside to outside, and to the acceleration (outermost curve). In the inset, the kurtosis $K(\tau) = \langle \delta_\tau v^4 \rangle / (\langle \delta_\tau v^2 \rangle)^2$ for the entire time interval $0.07\tau_\eta \div 2T_L$. The saturation of $K(\tau)$ at small time increments is an indication of the high numerical resolution.

come slightly sub-Gaussian for $\tau \approx T_L$, with a flatness $\langle (\delta_\tau v)^4 \rangle / (\langle \delta_\tau v^2 \rangle)^2 \approx 2.8$. Assuming a scaling law for the second-order Lagrangian velocity structure function in the time-range $\tau_\eta \ll \tau \ll T_L$, dimensional analysis [20], predicts $\langle (\delta_\tau v)^2 \rangle = C_0 \varepsilon \tau$. Remark that this prediction is not affected by intermittency corrections owing to its linear dependence on the energy dissipation rate [21, 22, 23]. Therefore, C_0 is expected to be universal. For our data the scaling behavior holds on a relatively narrow window in τ , leading to some uncertainty in the determination of the numerical prefactor. We estimate $C_0 = 5 \div 6$, not far from previous measurements [7, 24].

The intermittency in the PDFs of $\delta_\tau v$ can be conveniently quantified in terms of the Lagrangian velocity structure functions $S_p(\tau) = \langle (\delta_\tau v)^p \rangle$, that are expected to behave as power laws τ^{ξ_p} . The scaling exponents ξ_p can be evaluated by looking at the logarithmic slope $d \log S_p(\tau) / d \log \tau$ that should display a plateau in the range $\tau_\eta \ll \tau \ll T_L$. From Fig. 4 it is clear that it is very difficult to extract the values for ξ_p . However, by means of the extended self-similarity procedure [25], it is possible to estimate the relative exponents $\xi_4/\xi_2 = 1.7 \pm 0.05$, $\xi_5/\xi_2 = 2.0 \pm 0.05$, $\xi_6/\xi_2 = 2.2 \pm 0.07$, in fair agreement with those obtained in Ref. [7]. The range of time delays over which relative scaling occurs is $10\tau_\eta$ to $70\tau_\eta$. In this range anisotropic contributions induced by the large scale flow appear to influence the scaling properties.

It is interesting to remark that for values of τ in the range from τ_η to $10\tau_\eta$, the local slopes are significantly smaller and tend to accumulate around the value 2 for all orders. This relevant correction to scaling cannot be attributed to the influence of the dissipative range $\tau < \tau_\eta$,

since the latter would increase the value of the local slope, rather than decreasing it. A similar effect can be detected in Eulerian structure functions as well, yet the intensity in the latter case is much less pronounced. These strong deviations in the Lagrangian scaling laws are most likely due to the trapping events depicted in Fig. 2. Indeed, the long residence time within small-scale vortical structures introduces an additional weighting factor that enhances the effect with respect to Eulerian measurements. Further insight into the single particle dynamics can be




FIG. 4: Log-log plot of Lagrangian structure functions of orders $p = 2, 4, 6$ (bottom to top) vs τ . Bottom right: logarithmic local slopes $d \log S_p(\tau) / d \log \tau$ (same line styles). Top left: relative local slopes with respect to the second order structure function $d \log S_p(\tau) / d \log S_2(\tau)$, for $p = 4, 6$. Data refer to the v_x component. The two other velocity components exhibit slightly worse scaling due to anisotropy effects.

gained by investigating the various force terms concurring to determine the total particle acceleration

$$\frac{d^2 \mathbf{X}(t)}{dt^2} = -\nabla p(\mathbf{X}, t) + \nu \Delta \mathbf{u}(\mathbf{X}(t), t) + \mathbf{F}(\mathbf{X}(t), t).$$

As shown in the inset of Fig. 5, it is evident that the main source of acceleration is the pressure gradient [13], whereas the viscous term is important only near relatively strong acceleration events and the forcing contribution is negligible. A more quantitative test is given by the comparison of the PDFs of the three forces in Fig. 5. To conclude, we have presented the analysis of single-particle statistics in high Reynolds number flows. Our results fit well with previous experimental measurements. At variance with experiments, we can investigate the statistical properties of millions of particles on a wide range of time intervals, from a small fraction of the Kolmogorov time up to the integral correlation time. We found clear indications that velocity fluctuations along Lagrangian trajectories are affected by multiple-time dynamics. In the interval $10\tau_\eta < \tau < T_L$ we observed anomalous scaling for Lagrangian velocity structure functions. For frequencies of the order of τ_η^{-1} we noticed




FIG. 5: Log-log plot of PDFs for $-\partial_x p$, $\nu\Delta u_x$, F_x . The external forcing is virtually negligible, and the main contribution to large accelerations is made by pressure gradients. Inset: a typical evolution of the three terms along a particular trajectory. The strongest signal is $\partial_x p$ (dashed line), while the viscous force is activated only as a subleading response to pressure gradients (solid line). The force contribution is indistinguishable from zero.

that velocity fluctuations are affected by events where particles are trapped in vortex filaments. Events with trapping times much longer than expected on the basis of simple dimensional analysis appear frequently. The main novelty of Lagrangian single-particle velocity fluctuations with respect to Eulerian fluctuations is probably the capture of trajectories induced by small-scale structures. For example, the event analyzed in Fig. 2 would have a much smaller weight in an Eulerian analysis because of large-scale sweeping past the fixed probe. Among the most challenging open problems arising from our analysis are how to incorporate such dynamical processes in stochastic modelization of particle diffusion [3] and in the Lagrangian multifractal description [22, 23].

The simulations were performed within the keyproject “Lagrangian Turbulence” on the IBM-SP4 of Cineca (Bologna, Italy). We are grateful to C. Cavazzoni and G. Erbacci for resource allocation and precious technical assistance. We acknowledge support from EU under the contracts HPRN-CT-2002-00300 and HPRN-CT-2000-0162. We also thank E. L  v  que for useful dis-

cussions and B. Devenish for a careful reading of the manuscript.

-
- [1] S.B. Pope, *Annu. Rev. Fluid Mech.* **26**, 23 (1994).
 - [2] S.B. Pope, *Turbulent Flows*, Cambridge University Press (2000).
 - [3] B. Sawford, *Annu. Rev. Fluid Mech.* **33**, 289 (2001).
 - [4] P.K. Yeung, *Annu. Rev. Fluid Mech.* **34**, 115 (2002).
 - [5] A. La Porta, G.A. Voth, A.M. Crawford, J. Alexander and E. Bodenschatz, *Nature* **409**, 1017 (2001).
 - [6] G.A. Voth, A. La Porta, A. Crawford, E. Bodenschatz and J. Alexander, *J. Fluid Mech.* **469**, 121 (2002).
 - [7] N. Mordant, P. Metz, O. Michel and J.F. Pinton, *Phys. Rev. Lett.* **87**, 214501 (2001).
 - [8] S. Ott and J. Mann, *J. Fluid Mech.* **422**, 207 (2000).
 - [9] J.J. Riley and G.S. Patterson, Jr, *Phys. Fluids* **17**, 292 (1974).
 - [10] P.K. Yeung and S.B. Pope, *J. Fluid Mech.* **207**, 531 (1989).
 - [11] K.D. Squire and J.K. Eaton, *Phys. Fluids A* **3**, 130 (1991).
 - [12] P.K. Yeung, *J. Fluid Mech.* **427**, 241 (1991).
 - [13] P. Vedula and P.K. Yeung, *Phys. Fluids* **11**, 1208 (1999).
 - [14] G. Boffetta and I.M. Sokolov, *Phys. Rev. Lett.* **88**, 094501 (2002).
 - [15] T. Ishihara and Y. Kaneda, *Phys. Fluids* **14**, L69 (2002).
 - [16] L. Chevillard, S. G. Roux, E. Leveque, N. Mordant, J.F. Pinton and A. Arneodo, *Phys. Rev. Lett.* **91**, 214502 (2003).
 - [17] B.L. Sawford, P.K. Yeung, M.S. Borgas, P. Vedula, A. La Porta, A.M. Crawford and E. Bodenschatz, *Phys. Fluids* **15**, 3478 (2003).
 - [18] S. Chen, G. D. Doolen, R. H. Kraichnan and Z.-S. She, *Phys. Fluids A* **5**, 458 (1993).
 - [19] Y. Kaneda, T. Ishihara, M. Yokokawa, K. Itakura, and A. Uno, *Phys. Fluids* **15** L21 (2003).
 - [20] A. Monin and A. Yaglom, *Statistical Fluid Mechanics* MIT Press, Cambridge, MA (1975), Vol. 2.
 - [21] E.A. Novikov, *Phys. Fluids A* **1**, 326 (1989).
 - [22] M.S. Borgas, *Phil. Trans. R. Soc. Lond. A*, **342**, 379 (1993).
 - [23] G. Boffetta, F. De Lillo and S. Musacchio, *Phys. Rev. E* **66**, 066307 (2002).
 - [24] R.C. Lien and E.A. D’Asaro, *Phys. Fluids* **14**, 4456 (2002).
 - [25] R. Benzi, S. Ciliberto, R. Tripiccion, C. Baudet, F. Masaioli and S. Succi, *Phys. Rev. E* **48**, R29 (1993).